

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/367290372>

Bézier–Baskakov–Beta Type Operators

Article in *Filomat* · December 2022

DOI: 10.2298/FIL2219735K

CITATION

1

READS

169

3 authors, including:



Faruk Özger

Izmir Katip Celebi University

56 PUBLICATIONS 739 CITATIONS

[SEE PROFILE](#)



Jyoti Yadav

Central University of Haryana

4 PUBLICATIONS 2 CITATIONS

[SEE PROFILE](#)



Bézier-Baskakov-Beta Type Operators

Arun Kajla^a, Faruk Özger^b, Jyoti Yadav^a

^aDepartment of Mathematics, Central University of Haryana, Haryana 123031, India

^bDepartment of Engineering Sciences, İzmir Katip Çelebi University, 35620 İzmir, Türkiye

Abstract. In this study, we construct the Bézier-Baskakov-Beta type operators. We provide elements of Lipschitz type space, a direct approximation theorem by means of modulus of continuity $\omega_{\rho^{\ell}}(\zeta, t)$ ($0 \leq \ell \leq 1$) and approximation rate for functions having derivatives of bounded variation. We support the theoretical parts by computer graphics.

1. Introduction

Aral and Erbay [5] introduced Baskakov operators based on $\alpha \in [0, 1]$ as follows:

$$\mathcal{B}_m^{(\alpha)}(\zeta; x) = \sum_{j=0}^{\infty} b_{m,j}^{(\alpha)}(x) \zeta \left(\frac{j}{m} \right), \quad x \in [0, \infty). \quad (1)$$

Here

$$b_{m,j}^{(\alpha)}(x) = \frac{x^{j-1}}{(1+x)^{m+j-1}} \left[\frac{\alpha x}{(1+x)} \binom{m+j-1}{j} - (1-\alpha)(1+x) \binom{m+j-3}{j-2} + (1-\alpha)x \binom{m+j-1}{j} \right].$$

The operators $\mathcal{B}_m^{(\alpha)}(\zeta; x)$ reduce to Baskakov operators [6] for $\alpha = 1$.

Kajla et al. [19] considered a Durrmeyer type generalization of the operators (1) and gave the uniform convergence results. Gupta et al. [14] presented a family of approximation operators of exponential type and obtained some approximation theorem (See also [13]). In 2020, Mohiuddine et al. [21] Baskakov-Durrmeyer type operators based on the parameters and studied quantitative approximation results. Very recently, Mohiuddine et al. [20] introduced Stancu-Kantorovich form of the operators (1) and studied the direct results. Md. Nasiruzzaman et al. [22] constructed α -Schurer-Kantorovich operators and studied some direct results. N. L. Braha et al. [7] considered a Baskakov-Schurer-Szász-Stancu operators and established Korovkin type theorem, Grüss-Voronovskaya type theorem and the rate of convergence. H. M. Srivastava et al. in [26] proposed Szász-Mirakjan Beta-type operators and obtained the rate of convergence with the help of the classical and second moduli of continuity. Zeng and Chen [29] defined Bézier form of

2020 *Mathematics Subject Classification.* 26A15, 41A25, 41A35.

Keywords. Baskakov operators, Bézier operators, computer graphics, bounded variation.

Received: 21 January 2022; Revised: 17 June 2022; Accepted: 16 July 2022

Communicated by Hari M. Srivastava

Email addresses: arunkajla@cuh.ac.in (Arun Kajla), farukozger@gmail.com (Faruk Özger), jyoti201337@cuh.ac.in (Jyoti Yadav)

Bernstein-Durrmeyer operators and obtained the direct approximation theorem for functions of bounded variation. Srivastava and Gupta [25] obtained the convergence rate for Bézier form of BBH operators for functions of bounded variation. In 2007, Guo et al. [12] defined Baskakov-Bézier operators and studied the direct, inverse and equivalence approximation theorems with the help of Ditzian-Totik modulus of smoothness. Bézier type operators were studied by several researchers (cf. [1–3, 8, 15, 17, 27, 28, 31, 33]). Also uniform modulus of smoothness of certain operators were examined in [37–46].

Nasiruzzaman et al. [23] considered following Durrmeyer type generalization of the operators (1):

$$\mathcal{B}_{m,\alpha}^*(\zeta; x) = \sum_{j=0}^{\infty} b_{m,j}^{(\alpha)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} \zeta(t) dt, \tag{2}$$

where $B(j+1, m)$ is the beta function defined as

$$B(r, s) = \int_0^{\infty} \frac{w^{r-1}}{(1+w)^{r+s}} dw = \frac{\Gamma(r)\Gamma(s)}{\Gamma(r+s)}, \quad r, s > 0.$$

Nasiruzzaman et al. [23] studied uniform convergence of these operators in weighted spaces. A Durrmeyer type generalization which includes Beta function can be found in [4], and α type generalization of operators can be found in [35, 36].

Main goal of this article is to define the Bézier variant of the operators (2) and study some approximation properties e. g. the elements of Lipschitz type space, direct approximation theorem and the rate of approximation for functions having derivatives of bounded variation.

Let \mathfrak{L} denote the class of all Lebesgue measurable functions ζ on $[0, \infty)$ such that

$$\mathfrak{L} = \left\{ \zeta : \int_0^{\infty} \frac{|\zeta(t)|}{(1+t)^k} dt < \infty, \text{ for some positive integer } k \right\}.$$

For $\theta \geq 1$, we now consider the Bézier variant $\mathcal{B}_{m,\alpha,\theta}^*$ of the operators $\mathcal{B}_{m,\alpha}^*$ as follows:

$$\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) = \sum_{j=0}^{\infty} Q_{m,j,\alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} \zeta(t) dt, \tag{3}$$

where $Q_{m,j,\alpha}^{(\theta)}(x) = [V_{m,j,\alpha}(x)]^\theta - [V_{m,j+1,\alpha}(x)]^\theta$ with $V_{m,j,\alpha}(x) = \sum_{v=j}^{\infty} b_{m,v}^{(\alpha)}(x)$.

Alternatively we may rewrite the operators (3) as

$$\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) = \int_0^{\infty} U_{m,\alpha,\theta}(x, t) \zeta(t) dt, \quad x \in [0, \infty), \tag{4}$$

where

$$U_{m,\alpha,\theta}(x, t) = \sum_{j=0}^{\infty} Q_{m,j,\alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \frac{t^j}{(1+t)^{m+j+1}}.$$

For the sake of simplicity let $\bar{v} = (t-x)^2$ and $v = t-x$.

Lemma 1.1. [23] *Following relations are satisfied for $\mathcal{B}_{m,\alpha}^*$*

(a) $\mathcal{B}_{m,\alpha}^*(1; x) = 1;$

$$(b) \mathcal{B}_{m,\alpha}^*(t; x) = \frac{x(m + 2\alpha - 2)}{(m - 1)} + \frac{1}{(m - 1)}, \quad m > 1;$$

$$(c) \mathcal{B}_{m,\alpha}^*(t^2; x) = \frac{mx^2(m + 4\alpha - 3)}{(m - 2)(m - 1)} + \frac{x(4m + 10(\alpha - 1))}{(m - 2)(m - 1)} \frac{2}{(m - 2)(m - 1)}, \quad m > 2.$$

Lemma 1.2. [23] *Following relations are satisfied for $\mathcal{B}_{m,\alpha}^*$*

$$(a) \mathcal{B}_{m,\alpha}^*(v; x) = \frac{x(2\alpha - 1)}{(m - 1)} + \frac{1}{(m - 1)}, \quad m > 1;$$

$$(b) \mathcal{B}_{m,\alpha}^*(\bar{v}; x) = \frac{2x^2(m + 4\alpha - 3)}{(m - 2)(m - 1)} + \frac{2x(m + 5\alpha - 3)}{(m - 2)(m - 1)} + \frac{2}{(m - 2)(m - 1)}, \quad m > 2.$$

Corollary 1.3. *For $\lambda > 2$ and m sufficiently large, we have*

$$\mathcal{B}_{m,\alpha}^*(\bar{v}; x) \leq \frac{\lambda \rho^2(x)}{m}, \tag{5}$$

where $\rho(x) = \sqrt{x(1 + x)}$.

Lemma 1.4. *The inequality $\|\mathcal{B}_{m,\alpha}^*(\zeta)\| \leq \|\zeta\|$ is satisfied for $\zeta \in C_B[0, \infty)$.*

Lemma 1.5. *The inequality $\|\mathcal{B}_{m,\alpha,\theta}^*(\zeta)\| \leq \theta \|\zeta\|$ is satisfied for $\zeta \in C_B[0, \infty)$.*

Proof. Applying the inequality $|a^\beta - b^\beta| \leq \beta|a - b|$ with $0 \leq a, b \leq 1, \beta \geq 1$ and the definition of $Q_{m,j,\alpha}^{(\theta)}$, we obtain,

$$0 < [V_{m,j,\alpha}(x)]^\theta - [V_{m,j+1,\alpha}(x)]^\theta \leq \theta[V_{m,j,\alpha}(x) - V_{m,j+1,\alpha}(x)] = \theta b_{m,j}^{(\alpha)}(x). \tag{6}$$

From the operators $\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x)$ and Lemma 1.4, we get

$$\|\mathcal{B}_{m,\alpha,\theta}^*(\zeta)\| \leq \theta \|\mathcal{B}_{m,\alpha}^*(\zeta)\| \leq \theta \|\zeta\|.$$

□

2. Direct Theorem

From [30], we have

$$1 = V_{m,0,\alpha}(x) > V_{m,1,\alpha}(x) > \dots > V_{m,j,\alpha}(x) > V_{m,j+1,\alpha}(x) > \dots$$

$$0 < [V_{m,j,\alpha}(x)]^\theta - [V_{m,j+1,\alpha}(x)]^\theta \leq \theta b_{m,j}^{(\alpha)}(x), \quad \theta \geq 1. \tag{7}$$

We need following definitions to prove our results. Let $\rho(x) = \sqrt{x(1 + x)}, 0 \leq \ell \leq 1$, Ditizian-Totik modulus $\omega_{\rho^\ell}(\zeta, t)$ [9] is

$$\omega_{\rho^\ell}(\zeta, t) = \sup_{0 < h \leq t} \sup_{x \pm h\rho^\ell(x)/2 \geq 0} \left\{ \left| \zeta\left(x + \frac{h\rho^\ell(x)}{2}\right) - \zeta\left(x - \frac{h\rho^\ell(x)}{2}\right) \right| \right\},$$

and the appropriate Petree’s K -functional is defined by

$$\bar{K}_{\rho^\ell}(\zeta, t) = \inf_{g \in W_\ell} \{ \|\zeta - g\| + t\|\rho^\ell g'\| \}, \quad t > 0,$$

where $W_\ell = \{g : g \in AC_{loc}, \|\rho^\ell g'\| < \infty\}$. The relation $\bar{K}_{\rho^\ell}(\zeta, t) \sim \omega_{\rho^\ell}(\zeta, t)$ is well known [9, Theorem 3.1.2]. This implies there is a constant $M > 0$ such that

$$M^{-1}\omega_{\rho^\ell}(\zeta, t) \leq \bar{K}_{\rho^\ell}(\zeta, t) \leq M\omega_{\rho^\ell}(\zeta, t). \tag{8}$$

Lemma 2.1. For $\zeta \in W_\ell, \rho(x) = \sqrt{x(1+x)}, 0 \leq \ell \leq 1$ and $t, x > 0$, one has

$$\left| \int_x^t \zeta'(u) du \right| \leq 2^\ell \left(x^{-\ell/2} (1+t)^{-\ell/2} + \rho^{-\ell}(x) \right) |v| \|\rho^\ell \zeta'\|.$$

Proof. On an application of Hölder’s inequality, one has

$$\begin{aligned} \left| \int_x^t \zeta'(u) du \right| &\leq \|\rho^\ell \zeta'\| \left| \int_x^t \frac{du}{\rho^\ell(u)} \right| \\ &\leq \|\rho^\ell \zeta'\| |v|^{1-\ell} \left| \int_x^t \frac{du}{\rho(u)} \right|^\ell. \end{aligned}$$

Taking into account following two inequalities

$$\left| \int_x^t \frac{du}{\rho(u)} \right| \leq \left| \int_x^t \frac{du}{\sqrt{u}} \right| \left(\frac{1}{\sqrt{1+x}} + \frac{1}{\sqrt{1+t}} \right)$$

and

$$\left| \int_x^t \frac{du}{\sqrt{u}} \right| \leq \frac{2|v|}{\sqrt{x}}$$

one achieve following relations

$$\begin{aligned} \left| \int_x^t \zeta'(u) du \right| &\leq \|\rho^\ell \zeta'\| |v| \frac{2^\ell}{x^{\ell/2}} \left| \frac{1}{\sqrt{1+x}} + \frac{1}{\sqrt{1+t}} \right|^\ell \\ &\leq \|\rho^\ell \zeta'\| |v| \frac{2^\ell}{x^{\ell/2}} \left((1+t)^{-\ell/2} + (1+x)^{-\ell/2} \right) \end{aligned}$$

via inequality $|a + b|^p \leq |a|^p + |b|^p$ for $0 \leq p \leq 1$. \square

Lemma 2.2. Following inequality is satisfied

$$\mathcal{B}_{m,\alpha,\theta}^*((1+t)^{-k}; x) \leq C_k (1+x)^{-k} \tag{9}$$

for any non negative real number k , where C_k is a constant dependent on k .

Proof. For each $x \in [0, \infty)$, the result holds from (2). From (7), we have

$$\begin{aligned} \mathcal{B}_{m,\alpha,\theta}^*((1+t)^{-k}; x) &= \sum_{j=0}^{\infty} Q_{m,j,\alpha}^{(\theta)}(x) B(j+1, m) \int_0^\infty \frac{t^j}{(1+t)^{m+j+k+1}} dt \\ &= \sum_{j=0}^{\infty} \frac{Q_{m,j,\alpha}^{(\theta)}(x) \Gamma(m+j+1) \Gamma(m+k)}{\Gamma(m+k+j+1) \Gamma(m)} \\ &\leq \theta (1+x)^{-k} \frac{x^{j-1}}{(1+x)^{m+k+j-1}} \\ &\quad \times \left[\frac{\alpha x}{(1+x)} \binom{m+j-1}{j} - (1-\alpha)(1+x) \binom{m+j-3}{j-2} + (1-\alpha)x \binom{m+j-1}{j} \right] \\ &\quad \times \frac{\Gamma(m+j+1) \Gamma(m+k)}{\Gamma(m+k+j+1) \Gamma(m)}. \end{aligned} \tag{10}$$

Applying D’Alembert’s ratio test, the series on the right hand side (10) becomes convergent. This completes the proof. \square

For $c_1 \geq 0$ and $c_2 > 0$, Lipschitz-type space in two parameters [24] is given as

$$Lip_M^{(c_1, c_2)}(\vartheta) := \left\{ \zeta \in C_B[0, \infty) : |\zeta(t) - \zeta(x)| \leq M \frac{|v|^\vartheta}{(t + c_1x^2 + c_2x)^{\frac{\vartheta}{2}}}; x \in (0, \infty) \right\},$$

where M is a positive constant and $\vartheta \in (0, 1]$.

Theorem 2.3. Let $\zeta \in Lip_M^{(c_1, c_2)}(\vartheta)$. Then, one has

$$|\mathcal{B}_{m, \alpha, \theta}^*(\zeta; x) - \zeta(x)| \leq M \left(\theta \frac{\mu_{m, \alpha, 2}(x)}{(c_1x^2 + c_2x)} \right)^{\frac{\vartheta}{2}}$$

$\forall x > 0$, where $\mu_{m, \alpha, 2}(x) = \mathcal{B}_{m, \alpha, \theta}^*(\bar{v}; x)$.

Proof. For $\vartheta = 1$, one has

$$\begin{aligned} |\mathcal{B}_{m, \alpha, \theta}^*(\zeta; x) - \zeta(x)| &\leq \mathcal{B}_{m, \alpha, \theta}^*(|\zeta(t) - \zeta(x)|; x) \\ &\leq M \mathcal{B}_{m, \alpha, \theta}^* \left(\frac{|v|}{\sqrt{t + c_1x^2 + c_2x}}; x \right). \end{aligned} \tag{11}$$

Using the fact that $\frac{1}{\sqrt{t + c_1x^2 + c_2x}} < \frac{1}{\sqrt{c_1x^2 + c_2x}}$, the Cauchy-Schwarz inequality and applying Lemma 1.1 and (6), inequality (11) implies that

$$\begin{aligned} |\mathcal{B}_{m, \alpha, \theta}^*(\zeta; x) - \zeta(x)| &\leq M \frac{1}{\sqrt{c_1x^2 + c_2x}} \mathcal{B}_{m, \alpha, \theta}^*(|v|; x) \leq \frac{M}{\sqrt{c_1x^2 + c_2x}} (\mathcal{B}_{m, \alpha, \theta}^*(\bar{v}; x))^{1/2} \\ &\leq M \left(\sqrt{\theta \frac{\mu_{m, \alpha, 2}(x)}{c_1x^2 + c_2x}} \right). \end{aligned}$$

By the aid of Hölder inequality with $p = \frac{1}{\vartheta}$ and $q = \frac{1}{1-\vartheta}$, Lemma 1.1 and (6), one has

$$\begin{aligned} |\mathcal{B}_{m, \alpha, \theta}^*(\zeta; x) - \zeta(x)| &\leq \sum_{j=0}^{\infty} Q_{m, j, \alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} |\zeta(t) - \zeta(x)| dt \\ &\leq \left\{ \sum_{j=0}^{\infty} Q_{m, j, \alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \left(\int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} |\zeta(t) - \zeta(x)| dt \right)^{\frac{1}{\vartheta}} \right\}^{\vartheta} \\ &\leq \left\{ \sum_{j=0}^{\infty} Q_{m, j, \alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} |\zeta(t) - \zeta(x)|^{\frac{1}{\vartheta}} dt \right\}^{\vartheta} \\ &\leq M \left\{ \sum_{j=0}^{\infty} Q_{m, j, \alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} \frac{|v|}{\sqrt{t + c_1x^2 + c_2x}} dt \right\}^{\vartheta} \\ &\leq \frac{M}{(c_1x^2 + c_2x)^{\frac{\vartheta}{2}}} \left\{ \sum_{j=0}^{\infty} Q_{m, j, \alpha}^{(\theta)}(x) \frac{1}{B(j+1, m)} \int_0^{\infty} \frac{t^j}{(1+t)^{m+j+1}} |v| dt \right\}^{\vartheta} \\ &\leq \frac{M}{(c_1x^2 + c_2x)^{\frac{\vartheta}{2}}} (\mathcal{B}_{m, \alpha, \theta}^*(\bar{v}; x))^{\vartheta/2} \leq M \theta \frac{\mu_{m, \alpha, 2}^{\frac{\vartheta}{2}}(x)}{(c_1x^2 + c_2x)^{\frac{\vartheta}{2}}}, \end{aligned}$$

where $0 < \vartheta < 1$. \square

Next, we achieve a direct approximation theorem for Bézier-Baskakov-Beta type operators concerning Ditzian-Totik modulus of smoothness.

Theorem 2.4. Following inequality is satisfied for $\zeta \in C_B[0, \infty)$

$$|\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) - \zeta(x)| \leq C\omega_{\rho^\ell}\left(\zeta, \frac{\rho^{1-\ell}(x)}{\sqrt{m}}\right). \quad (12)$$

Proof. Bearing in mind definition of $\bar{K}_{\rho^\ell}(\zeta, t)$, for fixed m, x, ℓ one can take $g = g_{m,x,\ell} \in W_\ell$ so that

$$\|\zeta - g\| + \frac{\rho^{1-\ell}(x)}{\sqrt{m}} \|\rho^\ell g'\| \leq \bar{K}_{\rho^\ell}\left(\zeta, \frac{\rho^{1-\ell}(x)}{\sqrt{m}}\right). \quad (13)$$

Since $\mathcal{B}_{m,\alpha,\theta}^*(1; x) = 1$, one has

$$|\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) - \zeta(x)| \leq 2\|\zeta - g\| + |\mathcal{B}_{m,\alpha,\theta}^*(g; x) - g(x)|. \quad (14)$$

Taking into account Lemma 2.1 and the representation $g(t) = g(x) + \int_x^t g'(u)du$, we get

$$\begin{aligned} |\mathcal{B}_{m,\alpha,\theta}^*(g; x) - g(x)| &= \left| \mathcal{B}_{m,\alpha,\theta}^*\left(\int_x^t g'(u)du; x\right) \right| \\ &\leq 2^\ell \|\rho^\ell g'\| \left\{ \rho^{-\ell}(x) \mathcal{B}_{m,\alpha,\theta}^*(|v|; x) + x^{-\ell/2} \mathcal{B}_{m,\alpha,\theta}^*\left(\frac{|v|}{(1+t)^{\ell/2}}; x\right) \right\}. \end{aligned} \quad (15)$$

By the help of (7), Cauchy-Schwarz inequality and Corollary 1.3, we have

$$\begin{aligned} \mathcal{B}_{m,\alpha,\theta}^*(|v|; x) &\leq \left(\mathcal{B}_{m,\alpha,\theta}^*(\bar{v}; x)\right)^{1/2} \\ &\leq \frac{\sqrt{\theta\lambda}\rho(x)}{\sqrt{m}}. \end{aligned} \quad (16)$$

Similarly using Lemma 2.2, we get

$$\begin{aligned} \mathcal{B}_{m,\alpha,\theta}^*\left(\frac{|v|}{(1+t)^{\ell/2}}; x\right) &\leq \theta \mathcal{B}_{m,\alpha}^*\left(\frac{|v|}{(1+t)^{\ell/2}}; x\right) \\ &\leq \theta \left(\mathcal{B}_{m,\alpha}^*(\bar{v}; x)\right)^{1/2} \left(\mathcal{B}_{m,\alpha}^*((1+t)^{-\ell}; x)\right)^{1/2} \\ &\leq C_1 \frac{\sqrt{\theta\lambda}\rho(x)}{\sqrt{m}} (1+x)^{-\ell/2}. \end{aligned} \quad (17)$$

From (15)-(17), we get

$$|\mathcal{B}_{m,\alpha,\theta}^*(g; x) - g(x)| \leq C \|\rho^\ell g'\| \frac{\rho^{1-\ell}(x)}{\sqrt{m}}. \quad (18)$$

Using $\bar{K}_{\rho^\ell}(\zeta, t) \sim \omega_{\rho^\ell}(\zeta, t)$, (13), (14) and (18), we obtain (12). \square

3. Rate of Convergence

Suppose $\zeta \in DBV_\gamma(0, \infty)$, $\gamma \geq 0$, is the class of all functions defined on $(0, \infty)$, having a derivative of bounded variation on every finite subinterval of $(0, \infty)$, $\forall t > 0$ and $|\zeta(t)| \leq Mt^\gamma$.

Let $g(t)$ be a function of bounded variation on each finite subinterval of $(0, \infty)$. One has

$$\zeta(x) = \int_0^x g(t)dt + \zeta(0)$$

since $\zeta \in DBV_\gamma(0, \infty)$.

Lemma 3.1. *Following equalities are satisfied*

$$(a) \beta_{m,\alpha,\theta}(x, y) = \int_0^y U_{m,\alpha,\theta}(x, t) dt \leq \frac{\theta\lambda}{m} \frac{\rho^2(x)}{(x-y)^2}, \quad 0 \leq y < x,$$

$$(b) 1 - \beta_{m,\alpha,\theta}(x, z) = \int_z^\infty U_{m,\alpha,\theta}(x, t) dt \leq \frac{\theta\lambda}{m} \frac{\rho^2(x)}{(z-x)^2}, \quad x < z < \infty$$

for sufficiently large m , $x \in (0, \infty)$ and $\theta \geq 1, \lambda > 2$.

Proof. Taking into account (7) and Corollary 1.3, one has

$$\begin{aligned} \beta_{m,\alpha,\theta}(x, y) &= \int_0^y U_{m,\alpha,\theta}(x, t) dt \leq \int_0^y \left(\frac{x-t}{x-y}\right)^2 U_{m,\alpha,\theta}(x, t) dt \\ &\leq \mathcal{B}_{m,\alpha,\theta}^*(\bar{v}; x) (x-y)^{-2} \leq \theta \mathcal{B}_{m,\alpha}^*(\bar{v}; x) (x-y)^{-2} \\ &\leq \theta \frac{\lambda}{m} \frac{\rho^2(x)}{(x-y)^2} \end{aligned}$$

which completes proof of (a). Using the same manner one can prove part (b), so we omit this. \square

Theorem 3.2. *Suppose that $v_a^b(\zeta'_x)$ is the total variation of ζ'_x on $[a, b] \subset (0, \infty)$, $\lambda > 2$ and $\zeta \in DBV_\gamma(0, \infty)$, $\theta \geq 1$. Then, for every $x \in (0, \infty)$ and sufficiently large m , we have*

$$\begin{aligned} |\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) - \zeta(x)| &\leq \frac{\sqrt{\theta}}{\theta+1} \left| \zeta'(x+) + \theta \zeta'(x-) \right| \sqrt{\frac{\lambda}{m}} \rho(x) \\ &\quad + \sqrt{\frac{\lambda}{m}} \rho(x) \frac{\theta^{3/2}}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \\ &+ \frac{\theta\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_{x-x/j}^{x-x/j}(\zeta'_x) + \frac{x}{\sqrt{m}} v_{x-x/\sqrt{m}}^x(\zeta'_x) \\ &+ \frac{\theta\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_x^{x+x/j}(\zeta'_x) + \frac{x}{\sqrt{m}} v_x^{x+x/\sqrt{m}}(\zeta'_x). \end{aligned}$$

Here auxiliary function ζ'_x is defined as

$$\zeta'_x(t) = \begin{cases} \zeta'(t) - \zeta'(x-), & 0 \leq t < x, \\ 0, & t = x, \\ \zeta'(t) - \zeta'(x+) & x < t \leq 1. \end{cases}$$

Proof. Bearing in mind the result $\int_0^\infty U_{m,\alpha,\theta}(x, t) dt = \mathcal{B}_{m,\alpha,\theta}^*(1; x) = 1$, one has

$$\begin{aligned} \mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) - \zeta(x) &= \int_0^\infty [\zeta(t) - \zeta(x)] U_{m,\alpha,\theta}(x, t) dt \\ &= \int_0^\infty \left(\int_x^t \zeta'(u) du \right) U_{m,\alpha,\theta}(x, t) dt. \end{aligned} \tag{19}$$

By definition of function ζ'_x , for any $\zeta \in DBV_\gamma(0, \infty)$, one can have

$$\begin{aligned}\zeta'(t) &= \frac{1}{\theta+1} \left(\zeta'(x+) + \theta \zeta'(x-) \right) + \zeta'_x(t) \\ &\quad + \frac{1}{2} \left(\zeta'(x+) - \zeta'(x-) \right) \left(\operatorname{sgn}(v) + \frac{\theta-1}{\theta+1} \right) \\ &\quad + \delta_x(t) \left(\zeta'(x) - \frac{1}{2} \left(\zeta'(x+) + \zeta'(x-) \right) \right),\end{aligned}\tag{20}$$

where

$$\delta_x(t) = \begin{cases} 1, & x = t \\ 0, & x \neq t \end{cases}.$$

The following result is clear

$$\int_0^\infty U_{m,\alpha,\theta}(x,t) \int_x^t \left[\zeta'(x) - \frac{1}{2} \left(\zeta'(x+) + \zeta'(x-) \right) \right] \delta_x(u) du dt = 0.$$

By Cauchy-Schwarz inequality, (6) and simple computations, we have

$$\begin{aligned}E_1 &= \int_0^\infty \left(\int_x^t \frac{1}{\theta+1} \left(\zeta'(x+) + \theta \zeta'(x-) \right) du \right) U_{m,\alpha,\theta}(x,t) dt \\ &= \frac{1}{\theta+1} \left| \zeta'(x+) + \theta \zeta'(x-) \right| \int_0^\infty |v| U_{m,\alpha,\theta}(x,t) dt \\ &\leq \frac{1}{\theta+1} \left(\zeta'(x+) + \theta \zeta'(x-) \right) \left(\mathcal{B}_{m,\alpha,\theta}^*(\bar{v}; x) \right)^{1/2} \\ &\leq \frac{\sqrt{\theta}}{\theta+1} \left| \zeta'(x+) + \theta \zeta'(x-) \right| \sqrt{\frac{\lambda}{m}} \rho(x)\end{aligned}\tag{21}$$

and

$$\begin{aligned}E_2 &= \int_0^\infty \left(\int_x^t \frac{1}{2} \left(\zeta'(x+) - \zeta'(x-) \right) \left(\operatorname{sgn}(u-x) + \frac{\theta-1}{\theta+1} \right) du \right) U_{m,\alpha,\theta}(x,t) dt \\ &\leq \frac{\theta}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \int_0^\infty |v| U_{m,\alpha,\theta}(x,t) dt \\ &= \frac{\theta}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \mathcal{B}_{m,\alpha,\theta}^*(|v|; x) \\ &\leq \frac{\theta}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \left(\mathcal{B}_{m,\alpha,\theta}^*(\bar{v}; x) \right)^{1/2} \\ &\leq \frac{\theta^{3/2}}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \sqrt{\frac{\lambda}{m}} \rho(x).\end{aligned}\tag{22}$$

Taking into account (19)-(22), Lemma 1.2 and Corollary 1.3 following estimate is obtained

$$\begin{aligned}\left| \mathcal{B}_{m,\alpha,\theta}^*(\zeta; x) - \zeta(x) \right| &\leq \left| H_{m,\alpha,\theta}(\zeta'_x, x) + I_{m,\alpha,\theta}(\zeta'_x, x) \right| \\ &\quad + \frac{\sqrt{\theta}}{\theta+1} \left| \zeta'(x+) + \theta \zeta'(x-) \right| \sqrt{\frac{\lambda}{m}} \rho(x) \\ &\quad + \frac{\theta^{3/2}}{\theta+1} \left| \zeta'(x+) - \zeta'(x-) \right| \sqrt{\frac{\lambda}{m}} \rho(x),\end{aligned}\tag{23}$$

where

$$H_{m,\alpha,\theta}(\zeta'_x, x) = \int_0^x \left(\int_x^t \zeta'_x(u) du \right) U_{m,\alpha,\theta}(x, t) dt,$$

$$I_{m,\alpha,\theta}(\zeta'_x, x) = \int_x^\infty \left(\int_x^t \zeta'_x(u) du \right) U_{m,\alpha,\theta}(x, t) dt.$$

The rest of the proof is to estimate the terms $H_{m,\alpha,\theta}(\zeta'_x, x)$ and $I_{m,\alpha,\theta}(\zeta'_x, x)$. Following relations are satisfied

$$\begin{aligned} |H_{m,\alpha,\theta}(\zeta'_x, x)| &= \left| \int_0^x \left(\int_x^t \zeta'_x(u) du \right) d_t \beta_{m,\alpha,\theta}(x, t) \right| \\ &= \left| \int_0^x \beta_{m,\alpha,\theta}(x, t) \zeta'_x(t) dt \right| \\ &\leq \left(\int_0^y + \int_y^x \right) |\zeta'_x(t)| |\beta_{m,\alpha,\theta}(x, t)| dt \\ &\leq \theta \frac{\lambda \rho^2(x)}{m} \int_0^y v_i^x(\zeta'_x)(x-t)^{-2} dt + \int_y^x v_i^x(\zeta'_x) dt \\ &\leq \theta \frac{\lambda \rho^2(x)}{m} \int_0^y v_i^x(\zeta'_x)(x-t)^{-2} dt + \frac{x}{\sqrt{m}} v_{x-x/\sqrt{m}}^x(\zeta'_x) \end{aligned}$$

by the help of Lemma 3.1 with $y = x - x/\sqrt{m}$, the inequality $\int_a^b d_t \beta_{m,\alpha,\theta}(x, t) \leq 1$ for all $[a, b] \subseteq (0, \infty)$ and integration by parts.

Putting $u = x/(x - t)$ following relations are satisfied

$$\begin{aligned} \theta \frac{\lambda \rho^2(x)}{m} \int_0^{x-x/\sqrt{m}} (x-t)^{-2} v_i^x(\zeta'_x) dt &= \theta \frac{\lambda(1+x)}{m} \int_1^{\sqrt{m}} v_{x-x/u}^x(\zeta'_x) du \\ &\leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} \int_j^{j+1} v_{x-x/u}^x(\zeta'_x) du \\ &\leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_{x-x/j}^x(\zeta'_x). \end{aligned}$$

So the following inequality is satisfied

$$|H_{m,\alpha,\theta}(\zeta'_x, x)| \leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_{x-x/j}^x(\zeta'_x) + \frac{x}{\sqrt{m}} v_{x-x/\sqrt{m}}^x(\zeta'_x). \tag{24}$$

Applying Lemma 3.1 with $z = x + x/\sqrt{m}$ and via integration by parts, we have

$$\begin{aligned} |I_{m,\alpha,\theta}(\zeta'_x, x)| &= \left| \int_x^\infty \left(\int_x^t \zeta'_x(u) du \right) U_{m,\alpha,\theta}(x, t) dt \right| \\ &= \left| \int_x^z \left(\int_x^t \zeta'_x(u) du \right) d_t (1 - \beta_{m,\alpha,\theta}(x, t)) + \int_z^\infty \left(\int_x^t \zeta'_x(u) du \right) d_t (1 - \beta_{m,\alpha,\theta}(x, t)) \right| \end{aligned}$$

$$\begin{aligned}
 &= \left| \left[\left(\int_x^t \zeta'_x(u) du \right) (1 - \beta_{m,\alpha,\theta}(x,t)) \right]_x^z - \int_x^z \zeta'_x(t) (1 - \beta_{m,\alpha,\theta}(x,t)) dt \right. \\
 &\quad \left. + \int_z^\infty \left(\int_x^t \zeta'_x(u) du \right) d_t (1 - \beta_{m,\alpha,\theta}(x,t)) \right| \\
 &= \left| \left(\int_x^z \zeta'_x(u) du \right) (1 - \beta_{m,\alpha,\theta}(x,z)) - \int_x^z \zeta'_x(t) (1 - \beta_{m,\alpha,\theta}(x,t)) dt \right. \\
 &\quad \left. + \left[\left(\int_x^t \zeta'_x(u) du \right) (1 - \beta_{m,\alpha,\theta}(x,t)) \right]_z^\infty - \int_z^\infty \zeta'_x(t) (1 - \beta_{m,\alpha,\theta}(x,t)) dt \right| \\
 &= \left| \int_x^z \zeta'_x(t) (1 - \beta_{m,\alpha,\theta}(x,t)) dt + \int_z^\infty \zeta'_x(t) (1 - \beta_{m,\alpha,\theta}(x,t)) dt \right| \\
 &< \theta \frac{\lambda \rho^2(x)}{m} \int_z^\infty v_x^t (\zeta'_x(v))^{-2} dt + \int_x^z v_x^t (\zeta'_x) dt \\
 &\leq \theta \frac{\lambda \rho^2(x)}{m} \int_{x+x/\sqrt{m}}^\infty v_x^t (\zeta'_x(v))^{-2} dt + \frac{x}{\sqrt{m}} v_x^{x+x/\sqrt{m}} (\zeta'_x). \tag{25}
 \end{aligned}$$

Putting $u = x/v$ following relations are satisfied

$$\begin{aligned}
 \theta \frac{\lambda \rho^2(x)}{m} \int_{x+x/\sqrt{m}}^\infty v_x^t (\zeta'_x(v))^{-2} dt &= \theta \frac{\lambda \rho^2(x)}{xm} \int_0^{\sqrt{m}} v_x^{x+x/u} (\zeta'_x) du \\
 &\leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} \int_j^{j+1} v_x^{x+x/u} (\zeta'_x) du \\
 &\leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_x^{x+x/j} (\zeta'_x). \tag{26}
 \end{aligned}$$

Again combining (25)-(26), we get

$$|I_{m,\theta,\alpha}(\zeta'_x, x)| \leq \theta \frac{\lambda(1+x)}{m} \sum_{j=1}^{\lfloor \sqrt{m} \rfloor} v_x^{x+x/j} (\zeta'_x) + \frac{x}{\sqrt{m}} v_x^{x+x/\sqrt{m}} (\zeta'_x). \tag{27}$$

As a result, we get the required result combining (23), (24) and (27). □

4. Computer graphics

In this part, we provide some numerical results to see convergence of our operators. We first demonstrate behavior of polynomials $b_{m,j}^{(\alpha)}(x)$ for some j and α values, and $x \in [0, 5]$ in Figure 1 and Figure 2. We consider convergence and errors of convergence of our operators to the basic function

$$\zeta(x) = \sqrt{x}$$

in Figure 3 and Figure 4, respectively. We also consider

$$\zeta(x) = \frac{x+4}{x^2+6}$$

to see effectiveness of our operators. In Figure 5 (a), we choose $m = 20, \alpha = 0.5$ and $\theta = 2$ to see convergence of $\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x)$, where $x \in [0, 3]$. We select $m = 5, \alpha = 0.9$ and $\theta = 2$ in Figure 5 (b), $m = 10, \alpha = 0.8$ and $\theta = 3$ in Figure 5 (c), $m = 10, \alpha = 0.9$ and $\theta = 2$ in Figure 5 (d), $m = 15, \alpha = 0.9$ and $\theta = 2$ in Figure 5 (e) and $m = 25, \alpha = 0.5$ and $\theta = 2$ in Figure 5 (f) to observe approximation of $\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x)$ and corresponding errors. We can clearly say that, the operators $\mathcal{B}_{m,\alpha,\theta}^*(\zeta; x)$ have less errors even for small values of m .




Figure 1: Behavior of polynomials $b_{m,j}^{(\alpha)}(x)$ for $x \in [0, 5]$




Figure 2: Behavior of polynomials $b_{m,j}^{(\alpha)}(x)$ for some j and α values, and $x \in [0, 5]$




Figure 3: Approximation of operators for some m values




Figure 4: Errors of approximation for some m values




Figure 5: Convergence of operators and errors of approximation

Acknowledgements

The work of second author was supported by İzmir Katip Çelebi University Scientific Research Project Coordination Unit with project number 2020-GAP-MÜMF-0012.

References

- [1] U. Abel and V. Gupta, *An estimate of the rate of convergence of a Bézier variant of the Baskakov-Kantorovich operators for bounded variation functions*, Demonstratio Math. **36** (1) (2003) 123-136.
- [2] T. Acar and A. Kajla, *Blending type approximation by Bézier-summation-integral type operators*, Commun. Fac. Sci., Univ. Ank. Ser. A1 Math. Stat. **66** (2) (2018) 195–208.
- [3] P. N. Agrawal, N. Ispir and A. Kajla, *Approximation properties of Bézier-summation-integral type operators based on Polya-Bernstein functions*, Appl. Math. Comput. **259** (2015) 533-539.
- [4] A. Alotaibi, F. Özger, S.A. Mohiuddine and M. A. Alghamdi, *Approximation of functions by a class of Durrmeyer-Stancu type operators which includes Euler's beta function*. Adv. Differ. Equ. 2021, **13** (2021). <https://doi.org/10.1186/s13662-020-03164-0>
- [5] A. Aral and H. Erbay, *Parametric generalization of Baskakov operators*, Math. Commun. **24** (2019) 119-131.
- [6] V. A. Baskakov, *A sequence of linear positive operators in the space of continuous functions*, Dokl. Acad. Nauk. SSSR **113** (1957) 249-251.
- [7] N. L. Braha, T. Mansour, and H. M. Srivastava, *A parametric generalization of the Baskakov-Schurer-Szász-Stancu approximation operators*, Symmetry **13** (2021), Article ID 980, 1-24.
- [8] G. Chang, *Generalized Bernstein-Bézier polynomials*, J. Comput. Math. **1** (4) (1983) 322-327.
- [9] Z. Ditzian and V. Totik, *Moduli of Smoothness*, Springer, New York, (1987).
- [10] M. Goyal and P. N. Agrawal, *Bézier variant of the Jakimovski-Leviatan-Păltănea operators based on Appell polynomials*, Ann Univ Ferrara **63** (2017) 289-302
- [11] M. Goyal and P. N. Agrawal, *Bézier variant of the generalized Baskakov Kantorovich operators*, Boll. Unione Mat. Ital. **8** (2016), 229-238.
- [12] S. Guo, Q. Qi and G. Liu, *The central theorems for Baskakov-Bézier operators*, J. Approx. Theory **147** (2007) 112-124.
- [13] V. Gupta, A. M. Acu, and H. M. Srivastava, *Difference of some positive linear approximation operators for higher-order derivatives*, Symmetry **12** (2020) Article ID 915, 1-19
- [14] V. Gupta, M. López-Pellicer, and H. M. Srivastava, *Convergence estimates of a family of approximation operators of exponential type*, Filomat **34** (2020) 4329-4341.
- [15] V. Gupta, *On the Bézier variant of Kantorovich operators*, Comput. Math. Anal. **47** (2004) 227-232.
- [16] V. Gupta, *An estimate on the convergence of Baskakov-Bézier operators*. J. Math. Anal. Appl. **312** (2005) 280-288.
- [17] V. Gupta and R. P. Agarwal, *Convergence Estimates in Approximation Theory*, Springer, (2014).
- [18] A. Kajla, *On the Bézier variant of the Srivastava-Gupta operators*, Constr. Math. Anal. **1** (2018) 99-107.
- [19] A. Kajla, S. A. Mohiuddine, A. Alotaibi, M. Goyal and K. K. Singh, *Approximation by ϑ -Baskakov-Durrmeyer-type hybrid operators*, Iran. J. Sci. Technol. Trans. Sci. (2020) DOI 10.1007/s40995-020-00914-3.
- [20] S. A. Mohiuddine, N. Ahmad, F. Özger, A. Alotaibi and B. Hazarika, *Approximation by the parametric generalization of Baskakov-Kantorovich operators linking with Stancu operators*. Iran. J. Sci. Technol. Trans. A, Sci., **45** (2020) 593-605.
- [21] S. A. Mohiuddine, A. Kajla, M. Mursaleen, and M. A. Alghamdi, *Blending type approximation by τ -Baskakov-Durrmeyer type hybrid operators*, Adv. Differ. Equ. (2020) doi.org/10.1186/s13662-020-02925-1.
- [22] Md. Nasiruzzaman, H. M. Srivastava and S. A. Mohiuddine, *Approximation process based on parametric generalization of Schurer-Kantorovich operators and their bivariate form*, Proc. Nat. Acad. Sci. India Sect. A Phys. Sci. **92** (2022) 301-311.
- [23] Md. Nasiruzzaman, N. Rao, S. Wazir and R. Kumar, *Approximation on parametric extension of Baskakov-Durrmeyer operators on weighted spaces*. J. Inequal. Appl. **103** (2019) 1-11.
- [24] M. A. Özarslan and H. Aktuğlu, *Local approximation properties for certain King type operators*, Filomat **27** (1) 173-181.
- [25] H. M. Srivastava and V. Gupta, *Rate of convergence for the Bézier variant of the Bleimann-Butzer-Hahn operators*, Appl. Math. Lett. **18** (2005), 849–857
- [26] H. M. Srivastava, G. Ícoz, and B. Çekim, *Approximation properties of an extended family of the Szász-Mirakjan Beta-type operators*, Axioms **8** (2019) Article ID 111, 1-13.
- [27] H. M. Srivastava and X.M. Zeng, *Approximation by means of the Szász-Bézier integral operators*, International J. Pure Appl. Math. **14** (2004), No. 3, 283–294
- [28] X. M. Zeng, *On the rate of convergence of two Bernstein-Bézier type operators for bounded variation functions II*, J. Approx. Theory **104** (2000) 330-344.
- [29] X. M. Zeng and W. Chen, *On the rate of convergence of the generalized Durrmeyer type operators for functions of bounded variation*, J. Approx. Theory **102** (2000) 1-12.
- [30] X. M. Zeng and V. Gupta, *Rate of convergence of two Baskakov-Bézier type operators for locally bounded functions*, Comput. Math. Appl. **44** (2002) 1445-1453.
- [31] X. M. Zeng and A. Piriou, *On the rate of convergence of two Bernstein-Bézier type operators for bounded variation functions*, J. Approx. Theory **95** (1998) 369-387.
- [32] F. Özger, *Weighted statistical approximation properties of univariate and bivariate λ -Kantorovich operators*, Filomat **33** (11) (2019) 1-15.
- [33] F. Özger, *On new Bézier bases with Schurer polynomials and corresponding results in approximation theory*, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., **69** (1) (2020), 1-18.
- [34] H. M. Srivastava, F. Özger, S. A. Mohiuddine, *Construction of Stancu-type Bernstein operators Based on Bézier bases with shape parameter λ* , Symmetry **11** (3) (2019), Article 316.

- [35] S. A. Mohiuddine, F. Özger, *Approximation of functions by Stancu variant of Bernstein-Kantorovich operators based on shape parameter α* , Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM (2020) 114:70.
- [36] F. Özger, H. M. Srivastava, S. A. Mohiuddine, *Approximation of functions by a new class of generalized Bernstein-Schurer operators*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM (2020) 114:173.
- [37] K.J. Ansari, F. Özger, Z. Ödemiş Özger, *Numerical and theoretical approximation results for Schurer–Stancu operators with shape parameter λ* , Comp. Appl. Math. 41 (2022) 181.
- [38] M. Nasiruzzaman, A. Kiliçman, M. Ayman-Mursaleen, *Construction of q -Baskakov Operators by Wavelets and Approximation Properties*. Iran J Sci Technol Trans Sci 46, (2022) 1495–1503.
- [39] Q.B. Cai, K.J. Ansari, M. Temizer Ersoy, F. Özger, *Statistical blending-type approximation by a class of operators that includes shape parameters λ and α* , Mathematics 10(7) (2022) 1149.
- [40] U. Kadak, F. Özger, *A numerical comparative study of generalized Bernstein-Kantorovich operators*, Math. Found. Comput. 4(4) (2021), 311.
- [41] A. Kiliçman, M. Ayman-Mursaleen, M. Nasiruzzaman, *A note on the convergence of Phillips operators by the sequence of functions via q -calculus*, Demonstratio Mathematica, 55(1), (2022), 615-633.
- [42] F. Özger, E. Aljimi, M. Temizer Ersoy, *Rate of weighted statistical convergence for generalized blending-type Bernstein-Kantorovich operators*, Mathematics, 10(12) (2022), 2027.
- [43] F. Özger, K.J. Ansari, *Statistical convergence of bivariate generalized Bernstein operators via four-dimensional infinite matrices*, Filomat 36(2) (2022), 507–525.
- [44] H.M. Srivastava, E. Aljimi, B. Hazarika, *Statistical weighted $(N_\lambda, p, q)(E_\lambda, 1)$ A -summability with application to Korovkin's type approximation theorem*, Bull. Sci. Math. 178, (2022), 103146.
- [45] R. Aslan, *Some approximation results on λ -Szász-Mirakjan-Kantorovich operators*, Fundam. J. Math. Appl. 4(3) (2021), 150-158.
- [46] R. Aslan, *Approximation by Szász-Mirakjan-Durrmeyer operators based on shape parameter λ* , Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, (2022), 71.2: 407-421.