

5-6-2019

# Upper Record Values from Extended Exponential Distribution

Devendra Kumar

*Central University Haryana, Mahendergarh, India, devendrastats@gmail.com*

Sanku Dey

*St. Anthony's College, Shillong, India, sankud66@gmail.com*

Follow this and additional works at: <https://digitalcommons.wayne.edu/jmasm>

Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

## Recommended Citation

Kumar, D., & Dey, S. (2018). Upper record values from extended exponential distribution. *Journal of Modern Applied Statistical Methods*, 17(2), eP2687. doi: [10.22237/jmasm/1557149263](https://doi.org/10.22237/jmasm/1557149263)

This Regular Article is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Journal of Modern Applied Statistical Methods by an authorized editor of DigitalCommons@WayneState.

---

# Upper Record Values from Extended Exponential Distribution

## **Cover Page Footnote**

Acknowledgments The authors are grateful to the Editor and the anonymous referees for making many helpful comments and suggestions on an earlier version of this paper.

# Upper Record Values from Extended Exponential Distribution

**Devendra Kumar**

Central University of Haryana  
Mahendergarh, India

**Sanku Dey**

St. Anthony's College  
Shillong, India

---

Some recurrence relations are established for the single and product moments of upper record values for the extended exponential distribution by Nadarajah and Haghghi (2011) as an alternative to the gamma, Weibull, and the exponentiated exponential distributions. Recurrence relations for negative moments and quotient moments of upper record values are also obtained. Using relations of single moments and product moments, means, variances, and covariances of upper record values from samples of sizes up to 10 are tabulated for various values of the shape parameter and scale parameter. A characterization of this distribution based on conditional moments of record values is presented.

*Keywords:* Record values, single moment, product moment, recurrence relations, extended exponential distribution, conditional expectation, characterization

---

## Introduction

Record values are used in many statistical applications, statistical modeling and inference involving data pertaining to weather, athletic events, economics, life testing studies, and so on. Examples from *Guinness World Records* include the fastest time taken to recite the periodic table of the elements, shortest ever tennis matches both in terms of number of games and duration of time, or fastest indoor marathon. Several attempts are taken to make records, but records are made only when the attempt is a success. Usually, there is no data on all of the attempts made to break the record. Regarding the distributional properties of record-breaking data see Chandler (1952), Resnick (1973), Shorrock (1973), Glick (1978), Nevzorov (1987), Ahsanullah (1995), Balakrishnan and Ahsanullah (1993, 1994), Grunzień and Szyal (1997), and Arnold, Balakrishnan, and Nagaraja (1992, 1998).

---

The extended exponential has a decreasing probability function like an exponential distribution, but its mode at zero. It also allows for increasing, decreasing, and constant hazard rates like a Weibull distribution or an exponentiated exponential distribution. This distribution has explicit expressions of its survival function and failure rate function (see Nadarajah & Haghghi, 2011).

A random variable  $X$  is said to have an extended exponential distribution if its probability density function (pdf) is of the form

$$f(x; \alpha, \lambda) = \alpha \lambda (1 + \lambda x)^{\alpha-1} e^{-\lambda x}, \quad x > 0, \alpha, \lambda > 0, \quad (1)$$

and its corresponding cumulative distribution function (cdf) is

$$F(x; \alpha, \lambda) = 1 - e^{-\lambda x}, \quad x > 0, \alpha, \lambda > 0. \quad (2)$$

The hazard rate function is given by

$$h(x; \alpha, \lambda) = \alpha \lambda (1 + \lambda x)^{\alpha-1}, \quad x > 0, \alpha, \lambda > 0. \quad (3)$$

The properties of this distribution were studied by Nadarajah and Haghghi (2011). They obtained moment and maximum likelihood estimators of the distribution. This distribution is a particular member of the three-parameter generalized power Weibull distribution, introduced by Nikulin and Haghghi (2006). This distribution is a special case of the Gurvich, Dibenedetto, and Rande (1997) class as  $F(x) = 1 - \exp(-aG(x))$ , where  $G(x)$  is a monotonically increasing function of  $x$  with the only limitation  $G(x) \geq 0$ .

The concept of recurrence relations has its own importance. They are useful in reducing the number of operations necessary to obtain a general form for the function under consideration. Furthermore, they are used in characterizing distributions, which is an important area and permits the identification of the population distribution from the properties of the sample. Recurrence relations and identities have attained importance as they reduce the amount of direct computation, time, and labor. Recurrence relations for single and product moments of  $k$  record values from Weibull, Pareto, generalized Pareto, Burr, exponential, and Gumble distribution are derived by Pawlas and Szynal (1998, 1999, 2000). Kumar (2015, 2016), Kumar, Jain, and Gupta (2015), and Kumar, Kumar, Saran, and Jain (2017) established recurrence relations for moments of  $k^{\text{th}}$  record values for generalized

## UPPER RECORD VALUES

Rayleigh, Dagum, type-I generalized half logistic, and Kumaraswamy-Burr III distribution, respectively.

Some of the application areas of the extended exponential distribution include rainfall data analysis, earthquake frequency analysis, reliability, and survival analysis. In these and other application areas, the primary interest is in prediction of future events: what would be the magnitude of future rainfall, the magnitude of a future earthquake, etc. These predictions can be based on moments of record values.

### Record Values and Preliminaries

Let  $\{X_n, n \geq 1\}$  be a sequence of identically independently distributed (i.i.d.) random variables with cdf  $F(x)$  and pdf  $f(x)$ . The  $j^{\text{th}}$  order statistic of a sample  $(X_1, X_2, \dots, X_n)$  is denoted by  $X_{j:n}$ . For fixed  $k \geq 1$ , we define the sequence  $\{U^{(k)}(n), n \geq 1\}$  of  $k^{\text{th}}$  lower record times of  $X_1, X_2, \dots$  as follows:

$$U^{(k)}(1) = 1,$$

$$U^{(k)}(n+1) = \min \left\{ j > U^{(k)}(n), X_{j:j+k-1} > X_{U^{(k)}(n):U^{(k)}(n)+k-1} \right\}.$$

The sequences  $\{Y_n^{(k)}, n \geq 1\}$  with  $Y_n^{(k)} = X_{U^{(k)}(n):U^{(k)}(n)+k-1}$ ,  $n = 1, 2, \dots$ , are called the sequences of  $k^{\text{th}}$  upper record values of  $\{X_n, n \geq 1\}$ . For convenience, we shall also take  $Y_0^{(k)} = 0$ . Note that for  $k = 1$ , we have  $Y_1^{(1)} = X_{U^{(1)}(1)}$ ,  $n \geq 1$ , i.e. record values of  $\{X_n, n \geq 1\}$ . The joint pdf of  $k^{\text{th}}$  upper record values  $Y_1^{(k)}, Y_2^{(k)}, \mathbf{K}, Y_n^{(k)}$  can be given as the joint pdf of  $k^{\text{th}}$  upper record values of  $\{X_n, n \geq 1\}$  by

$$f_{Y_1^{(k)}, Y_2^{(k)}, \mathbf{K}, Y_n^{(k)}}(y_1, y_2, \mathbf{K}, y_n) = k^n \left( \prod_{i=1}^{n-1} \frac{f(y_i)}{\bar{F}(y_i)} \right) [\bar{F}(y_n)]^{k-1} f(y_n), \quad y_1 > y_2 > \mathbf{K} > y_n,$$

where  $\bar{F}(x) = 1 - F(x)$ . In view of above equation, the marginal pdf of  $X_n^{(k)}$ ,  $n \geq 1$ , is given by

$$f_{X_n^{(k)}}(x) = \frac{k^n}{(n-1)!} \left[ -\ln(\bar{F}(x)) \right]^{n-1} [\bar{F}(x)]^{k-1} f(x), \quad (4)$$

and the joint pdf of  $X_m^{(k)}$  and  $X_n^{(k)}$ ,  $1 \leq m < n$ ,  $n \geq 1$ , is given by

$$f_{X_m^{(k)}, X_n^{(k)}}(x, y) = \frac{k^n}{(m-1)!(n-m-1)!} [-\ln(\bar{F}(x))]^{m-1} \times [-\ln(\bar{F}(y)) + \ln(\bar{F}(y))]^{n-m-1} [\bar{F}(y)]^{k-1} \frac{f(x)}{\bar{F}(x)} f(y) \quad (5)$$

for  $x > y$ . Denote

$$\begin{aligned} \mu_{n:k}^{(r)} &= E\left(\left(X_n^{(k)}\right)^r\right) = \int_{-\infty}^{\infty} x^r f_{X_n^{(k)}}(x) dx, \quad r, n = 1, 2, K \\ \mu_{m,n:k}^{(r,s)} &= E\left(\left(X_m^{(k)}\right)^r \left(X_n^{(k)}\right)^s\right) = \int_{-\infty}^{\infty} \int_{-\infty}^x x^r y^s f_{X_m^{(k)}, X_n^{(k)}}(x, y) dy dx, \\ &1 \leq m \leq n-1, r, s = 1, 2, K \\ \mu_{m,n:k}^{(r,0)} &= E\left(\left(X_m^{(k)}\right)^r\right) = \mu_{m:k}^{(r)}, \quad r = 1, 2, K \\ \mu_{m,n:k}^{(0,s)} &= E\left(\left(X_n^{(k)}\right)^s\right) = \mu_{n:k}^{(s)}, \quad r = 1, 2, K \end{aligned}$$

## Relations for Single and Product Moments of Record Values

### Relations for Single Moments

For the extended exponential distribution in (1),

$$f(x; \alpha, \lambda) = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \lambda^{u+1} x^u \bar{F}(x; \alpha, \lambda). \quad (6)$$

The relation in (6) will be exploited to derive recurrence relations for the moments of record values for the extended exponential distribution. Establish the explicit expression for single moment of  $k^{\text{th}}$  record values  $E\left(\left(X_n^{(k)}\right)^r\right)$ . Using (4),

## UPPER RECORD VALUES

$$\mu_{n:k}^{(r)} = \frac{k^n}{(n-1)!} \int_0^\infty x^r [-\ln(\bar{F}(x))]^{n-1} [\bar{F}(x)]^{k-1} f(x) dx. \quad (7)$$

By setting  $t = -\ln[\bar{F}(x)]$  in (7) and simplifying,

$$\mu_{n:k}^{(r)} = \frac{k^n}{\lambda^r (n-1)!} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{(-1)^{p+r} \Gamma\left(\frac{p}{\alpha} + 1\right) \Gamma(r+1) \Gamma(n+q)}{k^q p! q! \Gamma\left(\frac{p}{\alpha} + 1 - q\right) \Gamma(r+1-q)}, \quad (8)$$

where  $\Gamma(a)$  denotes the complete gamma function defined by  $\Gamma(a) = \int_0^\infty t^{a-1} e^{-t} dt$ .

Specially, the first moment (mean) of the  $n^{\text{th}}$  record value is

$$\mu_n = \mu_n^{(1)} = \varphi(n, \alpha, \lambda, 1), \quad (9)$$

where

$$\varphi(n, \alpha, \lambda, a) = \frac{1}{\lambda^a (n-1)!} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{(-1)^{p+a} \Gamma\left(\frac{p}{\alpha} + 1\right) \Gamma(a+1) \Gamma(n+q)}{k^q p! q! \Gamma\left(\frac{p}{\alpha} + 1 - q\right) \Gamma(a+1-q)}.$$

In addition, the variance of  $U_n$  is

$$\sigma_n^2 = \mu_n^{(2)} - (\mu_n)^2 = \varphi(n, \alpha, \lambda, 2) - [\varphi(n, \alpha, \lambda, 1)]^2. \quad (10)$$

**Remark 1.** For  $k = 1$  in (8), deduce the explicit expression for single moments of upper record values for the extended exponential distribution.

Recurrence relations for single moments of  $k^{\text{th}}$  upper record values for cdf (2) are derived in the following theorem:

**Theorem 1.** For the extended exponential distribution given in (1) with fixed parameters  $\alpha, \lambda > 0, k, n = 1, 2, \dots$ , and the convention that  $\mu_{n:k}^{(0)} = 1$  and  $\mu_{0:k}^{(r)} = 0$ ,

$$\mu_{n:k}^{(r)} = \alpha k \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{\lambda^{u+1}}{(r+u+1)} \left[ \mu_{n:k}^{(r+u+1)} - \mu_{n-1:k}^{(r+u+1)} \right]. \quad (11)$$

**Proof.** For  $n \geq 1$  and  $r = 0, 1, 2, \dots$ , from (4),

$$\mu_{n:k}^{(r)} = \frac{k^n}{\alpha(n-1)!} \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \lambda^{u+1} \int_0^\infty [-\ln(\bar{F}(x))]^{n-1} [\bar{F}(x)]^k dx \quad (12)$$

Integrating by parts, using  $[-\ln(\bar{F}(x))]^{n-1}[\bar{F}(x)]^k$  for differentiation and the rest of the integrand for integration, we get the relation given in (11).

**Remark 2.** Setting  $k = 1$  in (11), we can deduce the recurrence relation for single moments of upper record values for the extended exponential distribution.

### Relations for Product Moments

Using (5), the explicit expression for the product moments of  $k^{\text{th}}$  upper record values  $\mu_{m,n:k}^{(r,s)}$  can be obtained as

$$\mu_{m,n:k}^{(r,s)} = \frac{k^n}{(m-1)!(n-m-1)!} \int_0^\infty x^r [-\ln(\bar{F}(x))]^{m-1} \frac{f(x)}{\bar{F}(x)} G(x) dx, \quad (13)$$

where

$$G(x) = \int_x^\infty y^s [-\ln(\bar{F}(y)) + \ln(\bar{F}(x))]^{n-m-1} [\bar{F}(y)]^{k-1} f(y) dy. \quad (14)$$

By setting  $w = -\ln(\bar{F}(y)) + \ln(\bar{F}(x))$  in (14), we obtain

$$\begin{aligned} & G(x) \\ &= \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{(-1)^{p(\frac{1}{\alpha}+1)+q+s} \Gamma\left(\frac{p}{\alpha}+1\right) \Gamma(s+1) \Gamma(n+q) [\bar{F}(x)]^k [-\ln \bar{F}(x)]^q \Gamma(n-m+q)}{k^{n-m+q} p! q! \Gamma\left(\frac{p}{\alpha}+1-q\right) \Gamma(s+1-p) \lambda^2} \end{aligned}$$

On substituting the above expression of  $G(x)$  into (13) and simplifying the resulting equation, obtain



## UPPER RECORD VALUES

$$\begin{aligned} \mu_{m,n;k}^{(r,s)} &= \frac{1}{\lambda^{s+r} (m-1)!(n-m-1)!} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{(-1)^{p(\frac{1}{\alpha}+1)+2q+s+r+u+m-1}}{p!q!u! \Gamma\left(\frac{p}{\alpha}+1-q\right)} \\ &\times \frac{\Gamma\left(\frac{p}{\alpha}+1\right) \Gamma(s+1) \Gamma(r+1) \Gamma(n-m+q)}{k^{q-m} p!q! \Gamma(s+1-p) \Gamma(r+1-u)} \frac{\partial^{m+q-1}}{\partial k^{m+q-1}} \mathbf{B}\left(\frac{u}{\lambda}+1, k\right) \end{aligned} \quad (15)$$

where  $\mathbf{B}(a, b)$  denotes the complete beta function defined by  $\mathbf{B}(a, b) = \Gamma(a)\Gamma(b) / \Gamma(a+b)$ . As a check, put  $s = 0$  in (15) and use (8),  $\mu_{m,n;k}^{(r,0)} = \mu_n^{(r)}$ .

For simplicity, we denote the  $(1, 1)^{\text{th}}$  moment of  $U_m$  and  $U_n$ , which are also called the simple product moment of these records, by  $\mu_{m,n}$ . The simple product moments are used for evaluating the covariances:

$$\sigma_{m,n} = \text{cov}(U_m, U_n) = \mu_{m,n} - \mu_m \mu_n.$$

**Remark 3.** Setting  $k = 1$  in (15), deduce the explicit expression for product moments of record values from the extended exponential distribution.

Making use of (6), drive recurrence relations for product moments of  $k^{\text{th}}$  upper record values:

**Theorem 2.** For  $1 \leq m \leq n-2$  and  $r, s = 0, 1, 2, \dots$ ,

$$\mu_{m,n;k}^{(r,s)} = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k \lambda^{u+1}}{(s+u+1)} \left[ \mu_{m,n;k}^{(r,s+1)} - \mu_{m,n-1;k}^{(r,s+1)} \right], \quad (16)$$

and for  $m \geq 2$  and  $r, s = 0, 1, 2, \dots$ ,

$$\mu_{m,m+1;k}^{(r,s)} = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k \lambda^{u+1}}{(s+u+1)} \left[ \mu_{m,m+1;k}^{(r,s+1)} - \mu_{m;k}^{(r,s+1)} \right]. \quad (17)$$

**Proof.** From equation (5) for  $1 \leq m \leq n-2$  and  $r, s = 0, 1, 2, \dots$ , on using (6),

$$\mu_{m,n;k}^{(r,s)} = \frac{k^n}{(m-1)!(n-m-1)!} \int_0^{\infty} x^r \left[ -\ln(\bar{F}(x)) \right]^{m-1} \frac{f(x)}{\bar{F}(x)} \mathbf{I}(x) dx \quad (18)$$

where

$$I(x) = \alpha \sum_{u=1}^{\alpha-1} \binom{\alpha-1}{u} \lambda^{u+1} \int_x^{\infty} y^{s+u} \left[ -\ln(\bar{F}(y)) + \ln(\bar{F}(x)) \right]^{n-m-1} [\bar{F}(y)]^k f(y) dy.$$

Integrating  $I(x)$  by parts,

$$\begin{aligned} I(x) &= -\alpha \sum_{u=1}^{\alpha-1} \binom{\alpha-1}{u} \frac{\lambda^{u+1}}{(s+u+1)} \\ &\quad \times \left\{ (n-m-1) \int_x^{\infty} y^{s+u+1} \left[ -\ln(\bar{F}(y)) + \ln(\bar{F}(x)) \right]^{n-m-2} [\bar{F}(y)]^{k-1} f(y) dy \right. \\ &\quad \left. + k \int_x^{\infty} y^{s+u+1} \left[ -\ln(\bar{F}(y)) + \ln(\bar{F}(x)) \right]^{n-m-1} [\bar{F}(y)]^{k-1} f(y) dy \right\} \end{aligned}$$

Substituting this expression in (18) and simplifying, it leads to (16). Proceeding in a similar manner for the case of  $n = m + 1$ , the recurrence relation (17) can be established.

**Remark 4.** Setting  $k = 1$  in (16) and (17), deduce the recurrence relation for the product moments of upper record values from the extended exponential distribution.

### Recurrence Relations for the Negative Moments

Let  $X_{U(1)}, X_{U(2)}, \dots$  be the sequence of the upper record values from the extended exponential distribution. The recurrence relation for the negative moments is given in the following theorem.

**Theorem 3.** For  $n \geq 1$  and  $s > 1$ ,

$$E\left(\frac{1}{X_{U(n)}^s}\right) = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k \lambda^{u+1}}{(u-s+1)} \left\{ E\left(\frac{1}{X_{U(n)}^{s-u-1}}\right) - E\left(\frac{1}{X_{U(n-1)}^{s-u-1}}\right) \right\}. \quad (19)$$

**Proof.** From (3), and for  $n \geq 1$  and  $s > 1$ ,

## UPPER RECORD VALUES

$$\begin{aligned}
 E\left(\frac{1}{X_{U(n)}^s}\right) &= \frac{k^n}{(n-1)!} \int_x^\infty x^{-s} \left[-\ln(\bar{F}(x))\right]^{n-1} [\bar{F}(x)]^{k-1} f(x) dx \\
 &= \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k^n \lambda^{u+1}}{(n-1)!} \int_x^\infty x^{u-s} \left[-\ln(\bar{F}(x))\right]^{n-1} [\bar{F}(x)]^k f(x) dx
 \end{aligned} \tag{20}$$

Integrating (20) by parts and simplifying the resultant expression, the result given in (19) is obtained.

**Remark 5.** Setting  $k = 1$  in (19), deduce the recurrence relation for the negative moments of upper record values from the extended exponential distribution.

### Recurrence Relations for the Quotient Moments

Let  $X_{U(1)}, X_{U(1)}, \dots$  be the sequence of the upper record values from the extended exponential distribution. Then the recurrence relation for the quotient moments is given by the following theorem:

**Theorem 5.** For  $1 \leq m \leq n$  and for  $r \geq 1$  and  $s \geq 1$ ,

$$E\left(\frac{X_{U(m;k)}^r}{X_{U(n;k)}^s}\right) = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k \lambda^{u+1}}{(u-s-1)} \left\{ E\left(\frac{X_{U(m;k)}^r}{X_{U(n)}^{s-u+1}}\right) - E\left(\frac{X_{U(m;k)}^r}{X_{U(n-1)}^{s-u+1}}\right) \right\}. \tag{21}$$

**Proof.** From (5), and for  $1 \leq m \leq n$ ,  $r \geq 1$ , and  $s \geq 1$ ,

$$\begin{aligned}
 E\left(\frac{X_{U(m;k)}^r}{X_{U(n;k)}^s}\right) &= \int_0^\infty \int_x^\infty \frac{x^r}{y^s} f_{X_m^{(k)}, X_n^{(k)}}(x, y) dy dx \\
 &= \frac{k^n}{(m-1)!(n-m-1)!} \int_0^\infty x^r \left[-\ln(\bar{F}(x))\right]^{m-1} \frac{f(x)}{\bar{F}(x)} h(x) dx
 \end{aligned} \tag{22}$$

where

$$h(x) = \int_x^\infty y^{-s} \left[-\ln(\bar{F}(y)) + \ln(\bar{F}(x))\right]^{n-m-1} [\bar{F}(y)]^k dy.$$

From the differential equation (6),

$$h(x) = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \lambda^{u+1} \int_x^{\infty} y^{u-s} \left[ -\ln(\bar{F}(y)) + \ln(\bar{F}(x)) \right]^{n-m-1} [\bar{F}(y)]^k dy. \quad (23)$$

Integrating (23) by parts and using (22), the result given in (21) is obtained.

**Corollary 1.** For  $m \geq 1$ ,  $r \geq 1$ , and  $s \geq 1$ ,

$$E\left(\frac{X_{U(m;k)}^r}{X_{U(m+1;k)}^s}\right) = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k\lambda^{u+1}}{(u-s-1)} \left\{ E\left(\frac{X_{U(m;k)}^r}{X_{U(m+1;k)}^{s-u+1}}\right) - E\left(X_{U(m;k)}^{(r+s-u-1)}\right) \right\}.$$

**Proof.** Upon substituting  $n = m + 1$  in (21) and simplifying, obtain the quotient moment (21).

**Remark 6.** Setting  $k = 1$  in (21), deduce the recurrence relation for the quotient moments of upper record values from the extended exponential distribution.

## Characterization

For upper record values, let  $L(a, b)$  stand for the space of all integrable functions on  $(a, b)$ . A sequence  $(h_n) \subset L(a, b)$  is called complete on  $L(a, b)$  if, for all functions  $g \in L(a, b)$ , the condition

$$\int_a^b g(x) h_n(x) dx = 0, \quad n \in \mathbf{N},$$

implies  $g(x) = 0$  almost everywhere (a.e.) on  $(a, b)$ . Start with the following result of Lin (1986, p. 595):

**Proposition.** Let  $n_0$  be any fixed non-negative integer,  $-\infty \leq a < b \leq \infty$ , and  $g(x) \geq 0$  an absolutely continuous function with  $g'(x) \neq 0$  a.e. on  $(a, b)$ . Then the sequence of functions  $\{(g(x))^n e^{-g(x)}, n \geq n_0\}$  is complete on  $L(a, b)$  iff  $g(x)$  is strictly monotone on  $(a, b)$ .

Using the above Proposition, a stronger version of Theorem 1 is obtained. Let  $\{X_n, n \geq 1\}$  be a sequence of i.i.d. continuous random variables with cdf  $F(x)$  and pdf  $f(x)$ . Let  $X_{L(n)}$  be the  $n^{\text{th}}$  upper record value; then the conditional pdf of  $X_{U(n)}$ , given  $X_{U(m)} = x$  and in view of (4) and (5), is

UPPER RECORD VALUES

$$f\left(X_{U(n)} \mid X_{U(m)} = x\right) = \frac{1}{(n-m-1)!} \left[-\ln \bar{F}(y) + \ln \bar{F}(x)\right]^{n-m-1} \frac{f(y)}{\bar{F}(x)}. \quad (24)$$

**Theorem 5.** Let  $X$  be an absolutely continuous random variable with cdf  $F(x)$  and pdf  $f(x)$  on the support  $(0, \infty)$ . Then, for  $m < n$ ,

$$E\left[X_{U(n)} \mid X_{U(m)} = x\right] = \frac{1}{(n-m-1)!} \sum_{p=0}^{\infty} \frac{\Gamma\left(\frac{1}{\alpha} + 1\right) \Gamma(n-m+p)}{p! \Gamma\left(\frac{1}{\alpha} + 1 - p\right) (1 + \lambda x)^{\alpha p - 1}} - 1 \quad (25)$$

if and only if  $F(x; \alpha, \lambda) = 1 - e^{-(1+\lambda x)^\alpha}$ ,  $x > 0$ ,  $\alpha, \lambda > 0$ .

**Proof.** From (24),

$$E\left[X_{U(n)} \mid X_{U(m)} = x\right] = \frac{1}{(n-m-1)!} \int_x^\infty y \left[ \ln \left( \frac{\bar{F}(x)}{\bar{F}(y)} \right) \right]^{n-m-1} \frac{f(y)}{\bar{F}(x)} dy. \quad (26)$$

By setting  $t = \ln(\bar{F}(x) / \bar{F}(y))$  from (2) in (26),

$$\begin{aligned} E\left[X_{U(n)} \mid X_{U(m)} = x\right] &= \frac{1}{(n-m-1)!} \sum_{p=0}^{\infty} \frac{\Gamma\left(\frac{1}{\alpha} + 1\right)}{p! \Gamma\left(\frac{1}{\alpha} + 1 - p\right) (1 + \lambda x)^{\alpha p - 1}} \\ &\quad \times \int_0^\infty t^{n-m+p-1} e^{-t} dt - \frac{1}{(n-m-1)!} \int_0^\infty t^{n-m+p-1} e^{-t} dt \end{aligned}$$

Simplifying the above expression, derive the relation given in (25). To prove the sufficient part, we have from (24) and (25)

$$\frac{1}{(n-m-1)!} \int_x^\infty y \left[-\ln \bar{F}(y) + \ln \bar{F}(x)\right]^{n-m-1} f(y) dy = \bar{F}(x) H_{n-m}(x), \quad (27)$$

where

$$H_{n-m}(x) = \frac{1}{(n-m-1)!} \sum_{p=0}^{\infty} \frac{\Gamma\left(\frac{1}{\alpha}+1\right)\Gamma(n-m+p)}{p!\Gamma\left(\frac{1}{\alpha}+1-p\right)(1+\lambda x)^{\alpha p-1}} - 1.$$

Differentiating (27) both sides with respect to  $x$ ,

$$\frac{f(x; \alpha, \lambda)}{\bar{F}(x; \alpha, \lambda)} = \frac{H'_{n-m}(x)}{[H_{n-m-1}(x) - H_{n-m}(x)]} = \alpha\lambda(1+\lambda x)^{\alpha-1},$$

which proves  $F(x; \alpha, \lambda) = 1 - e^{-(1+\lambda x)^\alpha}$ ,  $x > 0$ ,  $\alpha, \lambda > 0$ .

**Theorem 6.** Let  $X$  be a non-negative random variable having an absolutely continuous distribution function  $F(x)$  with  $F(0) = 0$  and  $0 < F(x) < 1$  for all  $x > 0$ . Then

$$\mu_{n:k}^{(r)} = \alpha k \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{\lambda^{u+1}}{(r+u+1)} [\mu_{n:k}^{(r+u+1)} - \mu_{n-1:k}^{(r+u+1)}] \quad (28)$$

if and only if  $F(x; \alpha, \lambda) = 1 - e^{-(1+\lambda x)^\alpha}$ ,  $x > 0$ ,  $\alpha, \lambda > 0$ .

**Proof.** The necessary part follows immediately from equation (11). However, if the recurrence relation in equation (28) is satisfied, then on using equation (4),

$$\begin{aligned} & \frac{k^n}{(n-1)!} \int_0^\infty x^r [\bar{F}(x)]^{k-1} [-\ln \bar{F}(x)]^{n-1} f(x) dx \\ &= -\alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k^{n+1} \lambda^{u+1}}{(r+u+1)(n-2)!} \\ & \quad \times \int_0^\infty x^{r+u+1} [\bar{F}(x)]^{k-1} [-\ln \bar{F}(x)]^{n-2} f(x) dx \\ &+ \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \frac{k^{n+1} \lambda^{u+1}}{(r+u+1)(n-1)!} \int_0^\infty x^{r+u+1} [\bar{F}(x)]^{k-1} [-\ln \bar{F}(x)]^{n-1} f(x) dx \end{aligned} \quad (29)$$

## UPPER RECORD VALUES

Integrating by parts, the first integral on the right-hand side of equation (29), and simplifying the resulting expression

$$\frac{f(x)}{\bar{F}(x)} = \alpha \sum_{u=0}^{\alpha-1} \binom{\alpha-1}{u} \lambda^{u+1} x^u,$$

which proves

$$F(x; \alpha, \lambda) = 1 - e^{-(1+\lambda x)^\alpha}, \quad x > 0, \alpha, \lambda > 0.$$

### Numerical Results

In Table 1, the values of means for  $n = 1, 2, \dots, 10$ ,  $\alpha = 1$  (1) 3, and  $\lambda = 0.5, 1.0$  are presented. Observe the means are decreasing with respect to  $n$  but increasing with respect to  $\alpha$ . In Table 2, the variances and covariances for different of values  $m$  and  $n$  and for  $\alpha = 1$  (1) 3 and  $\lambda = 0.5, 1.0$  are reported. Observe the variances and covariances are decreases with respect to both  $\alpha$  and  $\lambda$ .

**Table 1.** Means of record statistics from (8) for  $k = 1$

$n$	$\lambda = 0.5$			$\lambda = 1.0$		
	$\alpha = 1$	$\alpha = 2$	$\alpha = 3$	$\alpha = 1$	$\alpha = 2$	$\alpha = 3$
1	0.895563	1.791126	2.686689	0.352834	0.705668	1.058502
2	0.664068	1.328135	1.992203	0.199384	0.398769	0.598153
3	0.477761	0.955521	1.433282	0.107730	0.215459	0.323189
4	0.336661	0.673323	1.009984	0.056642	0.113284	0.169926
5	0.233775	0.467551	0.701326	0.029272	0.058543	0.087815
6	0.160622	0.321244	0.481866	0.014959	0.029918	0.044877
7	0.029140	0.219019	0.328529	0.007589	0.015178	0.022767
8	0.074238	0.148476	0.222713	0.003831	0.007663	0.011494
9	0.050114	0.100229	0.150343	0.001928	0.003856	0.005784
10	0.033724	0.067447	0.101171	0.000968	0.001936	0.002905

**Table 2.** Variances and covariances of record statistics

<i>m</i>	<i>n</i>	$\lambda = 0.5$			$\lambda = 1.0$		
		$\alpha = 1$	$\alpha = 2$	$\alpha = 3$	$\alpha = 1$	$\alpha = 2$	$\alpha = 3$
1	1	9.028346	8.315561	6.775821	3.648953	2.078890	1.693955
1	2	5.267103	4.930445	4.218403	2.055083	1.232611	1.054601
1	3	4.134393	3.916293	3.452603	1.060843	0.979073	0.863151
1	4	3.986392	3.479149	3.118780	0.965401	0.869787	0.779695
1	5	3.749502	3.261027	2.946530	0.957263	0.815257	0.736633
1	6	3.520278	3.143586	2.848129	0.893164	0.785897	0.712032
1	7	3.319361	3.077308	2.787888	0.820892	0.769327	0.696972
1	8	3.125013	3.038634	2.749184	0.812618	0.759659	0.687296
1	9	3.107148	3.015473	2.723474	0.806329	0.753868	0.680869
2	2	1.212775	1.591575	0.963569	0.303194	0.397894	0.240892
2	3	1.353388	1.035388	0.718258	0.338347	0.258847	0.179560
2	4	1.392593	0.864081	0.634816	0.348148	0.216020	0.158704
2	5	1.407665	0.790269	0.597443	0.351916	0.197567	0.149361
2	6	1.414247	0.754026	0.578157	0.353562	0.188506	0.144539
2	7	1.417306	0.735006	0.567199	0.354326	0.183751	0.141800
2	8	1.418774	0.724603	0.560513	0.354694	0.181151	0.140128
2	9	1.419492	0.718736	0.556216	0.354873	0.179684	0.139054
3	3	0.071090	0.084961	0.044993	0.284359	0.339844	0.179972
3	4	0.297832	0.258112	0.153716	0.074458	0.064528	0.038429
3	5	0.302410	0.228511	0.143493	0.075603	0.057128	0.035873
3	6	0.304325	0.215197	0.138739	0.076081	0.053799	0.034685
3	7	0.305199	0.208620	0.136260	0.076300	0.052155	0.034065
3	8	0.305616	0.205204	0.134846	0.076404	0.051301	0.033712
3	9	0.305819	0.203369	0.133981	0.076455	0.050842	0.033495
4	4	0.091612	0.083490	0.040061	0.022903	0.020873	0.010015
4	5	0.093130	0.071017	0.036969	0.023282	0.017754	0.009242
4	6	0.093731	0.065853	0.035655	0.023433	0.016463	0.008914
4	7	0.093999	0.063425	0.035024	0.023500	0.015856	0.008756
4	8	0.094126	0.062212	0.034689	0.023531	0.015553	0.008672
4	9	0.094187	0.061584	0.034490	0.023547	0.015396	0.008624
5	5	0.033398	0.022933	0.009890	0.008349	0.005733	0.002472
5	6	0.033594	0.020897	0.009503	0.008399	0.005224	0.002376
5	7	0.033680	0.019980	0.009330	0.008420	0.004995	0.002332
5	8	0.033720	0.019536	0.009244	0.008430	0.004884	0.002311
5	9	0.033739	0.019311	0.009198	0.008435	0.004828	0.002299
6	6	0.013079	0.006776	0.002589	0.003270	0.001694	0.000647
6	7	0.013107	0.006428	0.002539	0.003277	0.001607	0.000635
6	8	0.013120	0.006263	0.002516	0.003280	0.001566	0.000629
6	9	0.013126	0.006181	0.002504	0.003281	0.001545	0.000626
7	7	0.005369	0.002096	0.000701	0.001342	0.000524	0.000175
7	8	0.005373	0.002034	0.000694	0.001343	0.000509	0.000174
7	9	0.005369	0.002004	0.000691	0.001344	0.000501	0.000173
8	8	0.002278	0.000667	0.000193	0.000569	0.000167	0.000048
8	9	0.002258	0.000656	0.000183	0.000570	0.000164	0.000043



## Conclusion

The upper record values from the extended exponential model were considered, and exact explicit expressions were obtained, as well as recurrence relations for the single and product moments of record values. Recurrence relations for negative moments and quotient moments of upper record values were also obtained. The recurrence relations obtained in the paper allows us to evaluate the means, variances, and covariances of all upper record values for all sample sizes in a simple recursive manner.

## Acknowledgements

The authors are grateful to the Editor and the anonymous referees for making many helpful comments and suggestions on an earlier version of this paper.

## References

- Ahsanullah, M. (1995). *Record statistics*. New York, NY: Nova Science Publishers.
- Arnold, B. C., Balakrishnan, N., & Nagaraja, H. N. (1992). *A first course in order statistics*. New York, NY: John Wiley and Sons.
- Arnold, B. C., Balakrishnan, N., & Nagaraja, H. N. (1998). *Records*. New York, NY: John Wiley and Sons. doi: [10.1002/9781118150412](https://doi.org/10.1002/9781118150412)
- Balakrishnan, N., & Ahsanullah, M. (1993). Relations for single and product moments of record values from exponential distribution. *Journal of Applied Statistical Science*, 2(1) 73-87.
- Balakrishnan, N., & Ahsanullah, M. (1994). Recurrence relations for single and product moments of record values from generalized Pareto distribution. *Communications in Statistics – Theory and Methods*, 23(10), 2841-2852. doi: [10.1080/03610929408831419](https://doi.org/10.1080/03610929408831419)
- Chandler, K. N. (1952). The distribution and frequency of record values. *Journal of the Royal Statistical Society: Series B (Methodological)*, 14(2), 220-228. doi: [10.1111/j.2517-6161.1952.tb00115.x](https://doi.org/10.1111/j.2517-6161.1952.tb00115.x)
- Glick, N. (1978). Breaking records and breaking boards. *The American Mathematical Monthly*, 85(1), 2-26. doi: [10.1080/00029890.1978.11994501](https://doi.org/10.1080/00029890.1978.11994501)

Grunzień, Z., & Szynal, D. (1997). Characterization of uniform and exponential distributions via moments of the  $k$ th record values with random indices. *Journal of Applied Statistical Science*, 5(4), 259-266.

Gurvich, M. R., Dibenedetto, A. T., & Rande, S. V. (1997). A new statistical distribution for characterizing the random strength of brittle materials. *Journal of Materials Science*, 32(10), 2559-2564. doi: 10.1023/a:1018594215963

Kumar, D. (2015). Explicit expressions and statistical inference of generalized Rayleigh distribution based on lower record values. *Mathematical Methods of Statistics*, 24(3), 225-241. doi: 10.3103/s1066530715030035

Kumar, D. (2016).  $K$ -th lower record values from of Dagum distribution. *Discussions Mathematicae Probability and Statistics*, 36, 25-41. doi: 10.7151/dmps.1183

Kumar, D., Jain, N., & Gupta, S. (2015). The type I generalized half-logistic distribution based on upper record values. *Journal of Probability and Statistics*, 2015, Article ID 393608. doi: 10.1155/2015/393608

Kumar, D., Kumar, M., Saran, J., & Jain, N. (2017). The Kumaraswamy-Burr III distribution based on upper record values. *American Journal of Mathematical and Management Sciences*, 36(3), 205-228. doi: 10.1080/01966324.2017.1318099

Lin, G. D. (1986). On a moment problem. *Tohoku Math Journal*, 38(4), 595-598. doi: 10.2748/tmj/1178228411

Nadarajah, S., & Haghighi, F. (2011). An extension of the exponential distribution. *Statistics*, 45(6), 543-558. doi: 10.1080/02331881003678678

Nevzorov, V. B. (1987). Records. *Theory of Probability & Its Applications*, 32(2), 201-228. doi: 10.1137/1132032

Nikulin, M., & Haghighi, F. (2006). A chi-squared test for the generalized power Weibull family for the head-and-neck cancer censored data. *Journal of Mathematical Sciences*, 133(3), 1333-1341. doi: 10.1007/s10958-006-0043-8

Pawlas, P., & Szynal, D. (1998). Relations for single and product moments of  $k$ -th record values from exponential and Gumbel distributions. *Journal of Applied Statistical Science*, 7(1), 53-61.

Pawlas, P., & Szynal, D. (1999). Recurrence relations for single and product moments of  $k$ -th record values from Pareto, generalized Pareto and Burr distributions. *Communications in Statistics – Theory and Methods*, 28(7), 1699-1709. doi: 10.1080/03610929908832380

## UPPER RECORD VALUES

Pawlas, P., & Szynal, D. (2000). Recurrence relations for single and product moments of k-th record values from Weibull distribution and a characterization. *Journal of Applied Statistical Science*, 10(1), 17-25.

Resnick, S. I. (1973). *Extreme values, regular variation and point processes*. New York, NY: Springer-Verlag.

Shorrock, R. W. (1973). Record values and inter-record times. *Journal of Applied Probability*, 10(3), 543-555. doi: 10.2307/3212775